



## **Speciation of Heavy Metals - an important parameter for risk assessment of feed and food safety in aquaculture**

**Sloth, Jens Jørgen**

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Sloth, J. J. (Author). (2011). Speciation of Heavy Metals - an important parameter for risk assessment of feed and food safety in aquaculture. Sound/Visual production (digital)  
<https://www.was.org/WasMeetings/meetings/Default.aspx?code=WA2011>

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# SPECIATION OF HEAVY METALS

– AN IMPORTANT PARAMETER FOR RISK ASSESSMENT OF FEED AND  
FOOD SAFETY IN AQUACULTURE

**Jens J. Sloth**

*National Food Institute  
Technical University of Denmark*



[www.confidence.eu](http://www.confidence.eu)



# Current situation in EU legislation:



## Foodstuffs

MLs for Pb, Cd, Hg and Sn  
EU directive 2006/1881/EC



## Animal feedingstuffs

MLs for As, Pb, Cd and Hg  
EU directive 2002/32/EC


**Only maximum levels for  
total concentration of the metals**



# Speciation of elements



Novel speciation alternative – SPE-HG-AAS



The analytical speciation workhorse – HPLC-ICPMS




HPLC (GC)



Column



ICPMS



Chromatogram

Example Hg speciation




# Selected sample types

## ➤ Focus on marine feed and seafood

Seafood is the main dietary source of **arsenic** and mercury

Average EU: ~1 µg/kg bw/day



Fish and seafood →



EFSA (2009), Scientific Opinion on Arsenic in Food






# Selected sample types

## ➤ Focus on marine feed and seafood


Seafood is the main dietary source of arsenic and **mercury**






Ref: Danish Food Adm, 2004.




# Arsenic compounds in the marine environment



More than **50** different arsenic species have been found in the marine environment




# Arsenic compounds in the marine environment




Arsenocosis

**Most toxic form of arsenic!!**


## Inorganic arsenic




Arsenous acid  
As(III)



Arsenic acid  
As(V)




Trimethylarsonioriboside












# Arsenic compounds in the marine environment



AB="fish arsenic"  
(Chapman, Analyst, 1920)


## Organoarsenic compounds - trimethylated species



Trimethylarsoniopropionic acid  
**TMAP**

Trimethylarsonioriboside

**Predominant form  
of arsenic in most  
seafood!!**




# Arsenic compounds in the marine environment




**Predominant form of arsenic in algae!!**

## Arsenosugars











> 15 different identified



Dimethylarsinoylribosides



Trimethylarsonioribosides



# Commission directive 2009/114/EC (amendment)

Undesirable substances	Products intended for animal feed	Maximum content in mg/kg (ppm) relative to a feedingstuff with a moisture content of 12 %
(1)	(2)	(3)
1. Arsenic (*) (**)	Feed materials with the exception of:	2
	— meal made from grass, from dried lucerne and from dried clover, and dried sugar beet pulp and dried molasses sugar beet pulp,	4
	— palm kernel expeller,	4 (***)
	— phosphates and calcareous marine algae,	10
	— calcium carbonate,	15
	— magnesium oxide,	20
	— feedingstuffs obtained from the processing of fish or other marine animals, including fish,	25 (***)
	— seaweed meal and feed materials derived from seaweed,	40 (***)
	Iron particles used as tracer.	50
	Additives belonging to the functional group of compounds of trace elements except:	30



Only max levels for total arsenic!!

## FOOTNOTE

(\*\*\*) Upon request of the competent authorities, the responsible operator must perform an analysis to demonstrate that the content of inorganic arsenic is lower than 2 ppm in the following products: (a) zinc oxide, manganese oxide and copper oxide, (b) complete feedingstuffs with the exception of: (i) complete feedingstuffs for fish and complete feedingstuffs for fur animals, (c) complementary feedingstuffs with the exception of: (i) mineral feedingstuffs.

Complete feedingstuffs with the exception of:	2
— complete feedingstuffs for fish and complete feedingstuffs for fur animals,	10 (***)
Complementary feedingstuffs with the exception of:	4
— mineral feedingstuffs,	12

Speciation analysis is required !!




# SPE-HG-AAS – a novel speciation alternative...

$\mu$ -wave extraction


Separation by SPE

Detection by HG-AAS

Inexpensive detection system



Sequential elution for selective off-line separation of inorg As from organo As species by SPE



HPLC-ICPMS of SPE fractions


# Performance characteristics from in-house validation

μ-wave  
extraction

Separation by  
SPE

Detection by  
HG-AAS

Parameter	Result
Analysis time	2 x 7 h for 24 samples
LoD (mg/kg)	0.08
LoQ (mg/kg)	0.16
Repeatability (%RSD)	3 - 7
Accuracy (%)	90 - 104



Further details on  
poster and handouts



# EFSA (2009) and JECFA (2010) opinions on arsenic in food

- Old PTWI value (WHO, 1988) was withdrawn
- **NEW!**  $\text{BMDL}_{1.0} = 0.3 - 8 \mu\text{g/kg bw per day}$  for inorganic arsenic
- => EU dietary exposures within this range
- => Risk to some consumers cannot be excluded



- **NEW!**  $\text{BMDL}_{0.5} = \underline{3 \mu\text{g/kg bw per day}}$  for inorganic arsenic
- => *0.5% increased incidence of lung cancer for 12 y exposure*




- "...more accurate information on the inorganic arsenic content of foods is needed to improve assessments of dietary exposures to inorganic arsenic"
- "...need for validated methods for selective determination of inorganic arsenic in food matrices"





# Speciation analysis of Mercury by HPLC-ICPMS




MeHg is a priority  
contaminant

- Adverse effects on the central nervous system
- Affects the learning ability of children
- Bioaccumulation and biomagnification in the aquatic food chain
- Biomethylation in the food chain:  $\text{iHg} \rightarrow \text{MeHg}$
- Toxicity:  $\text{MeHg} > \text{inorganic Hg}$
- PTWI MeHg:  $1.6 \mu\text{g/kg bw/week}$



# Speciation analysis of Mercury by HPLC-ICPMS



*HPLC-ICPMS chromatogram of DORM-3 (Dogfish muscle)*

CRM	Certified (mg/kg)	Result (mg/kg)
DORM-2 (dogfish muscle)	4.47 +/- 0.32	4.21
DORM-3 (dogfish muscle)	0.355 +/- 0.056	0.35
TORT-2 (Lobster hepatopancreas)	0.152 +/- 0.013	0.16



# Conclusion

- Total concentration info – not always sufficient!
- Speciation analysis for improved risk assessment
- Need for speciation methods for future monitoring
- ....and in feed and food control



## ***Thanks for your attention!***

Further information:

**Speciation – chemical analysis:** [jjsl@food.dtu.dk](mailto:jjsl@food.dtu.dk) (Jens J. Sloth)



Miscellaneous info on speciation: EVISA homepage; [www.speciation.net](http://www.speciation.net)  **evisa.**

Further developments in **CONFIDENCE**: [www.confidence.eu](http://www.confidence.eu)

